

Induced structures on product of spheres

Cristina-Elena Hreţcanu

Abstract

The purpose of this paper is to give an effective construction for some induced structures on spheres or product of spheres of codimension 1, 2 and 3, respectively, in Euclidean space endowed with an almost product structure.

1 Introduction

Let $(\widetilde{M}, \widetilde{g})$ be a Riemannian manifold equipped with a Riemannian metric tensor $\widetilde{g}$ and a $(1,1)$ tensor field $\widetilde{P}$ such that

$$(1.1) \quad \widetilde{P}^2 = \epsilon I,$$

where I is the identity on $\widetilde{M}$ and $\epsilon = \pm 1$. We suppose that $\widetilde{g}$ and $\widetilde{P}$ are compatible in the sense that we have

$$(1.2) \quad \widetilde{g}(\widetilde{P}U, \widetilde{P}V) = \widetilde{g}(U, V),$$

which is equivalent to

$$(1.3) \quad \widetilde{g}(\widetilde{P}U, V) = \epsilon \widetilde{g}(U, \widetilde{P}V),$$

for each $U, V \in \chi(\widetilde{M})$, where $\chi(\widetilde{M})$ is the Lie algebra of the vector fields on $\widetilde{M}$.

For $\epsilon = 1$ we obtain that $\widetilde{P}$ is an almost product structure and the Riemannian manifold $(\widetilde{M}, \widetilde{g}, \widetilde{P})$, with the compatibility relation (1.2), becomes an almost product Riemannian manifold.

Let M be a n -dimensional submanifold of codimension r ($n, r \in \mathbb{N}^*$) in a Riemannian manifold $(\widetilde{M}, \widetilde{g}, \widetilde{P})$ which satisfied the relations (1.1) and (1.2).

We note the tangent space of M in a point $x \in M$ by $T_x(M)$ and the normal space of M in $\widetilde{M}$ by $T_x^\perp(M)$. For $\alpha \in \{1, \dots, r\}$, let $(N_1, \dots, N_r) := (N_\alpha)$ be an orthonormal basis in $T_x^\perp(M)$, for every $x \in M$.

In the same manner like in [1] and [7], we construct the $(a, \epsilon)f$ Riemannian structure, which is a structure $(P, g, \epsilon \xi_\alpha, u_\alpha, (a_{\alpha\beta})_r)$, induced on a submanifold M in a Riemannian manifold $(\widetilde{M}, \widetilde{g}, \widetilde{P})$ ($\widetilde{P}$ and $\widetilde{g}$ verify the

relation (1.1) and (1.2)), where P is an $(1,1)$ -tensor field on M , g is the induced metric on submanifold M , ξ_α are tangent vector fields on M , u_α are 1-forms on M and $(a_{\alpha\beta})_r$ is a $r \times r$ matrix of real functions on M .

The $(a, \epsilon)f$ Riemannian structure induced on M by the $\tilde{P}$ structure on $(\tilde{M}, \tilde{g})$ (which verifies the relations (1.1) and (1.2)) is a generalization of the almost r -paracontact Riemannian structure([3]). The case of the $(a, -1)f$ Riemannian structure was studied by K. Yano and M. Okumura (in [17] and [18]).

2 $(a, \epsilon)f$ Riemannian structure

Let M be a n -dimensional submanifold of codimension r ($n, r \in \mathbb{N}^*$) in a Riemannian manifold $(\tilde{M}, \tilde{g})$, equipped by an $(1,1)$ -tensor field $\tilde{P}$, such that $\tilde{g}$ and $\tilde{P}$ verify the conditions (1.1) and (1.2). The decomposition of vector fields $\tilde{P}X$ and $\tilde{P}N_\alpha$, respectively, in the tangential and normal components on the submanifold M in $\tilde{M}$ has the form:

$$(2.1) \quad \tilde{P}X = PX + \sum_{\alpha=1}^r u_\alpha(X)N_\alpha,$$

for every $X \in \chi(M)$ and

$$(2.2) \quad \tilde{P}N_\alpha = \epsilon \xi_\alpha + \sum_{\beta=1}^r a_{\alpha\beta} N_\beta, \quad (\epsilon = \pm 1)$$

for every $\alpha \in \{1, \dots, r\}$.

We use the notation $a := (a_{\alpha\beta})_r$ for the matrix which result from the relation (2.2).

Definition 2.1. *An $(a, \epsilon)f$ Riemannian structure on a n -dimensional submanifold M of codimension r ($n, r \in \mathbb{N}^*$) in a Riemannian manifold $(\tilde{M}, \tilde{g})$ is a structure $(P, g, \epsilon \xi_\alpha, u_\alpha, (a_{\alpha\beta})_r)$ induced on M by the structure $(\tilde{P}, \tilde{g})$ from $\tilde{M}$ (where $\tilde{g}$ and $\tilde{P}$ verify the conditions (1.1) and (1.2)).*

Theorem 2.1. ([7]) *Let M be a n -dimensional submanifold of codimension r ($n, r \in \mathbb{N}^*$) in a Riemannian manifold $(\tilde{M}, \tilde{g})$, equipped by an $(1,1)$ -tensor field $\tilde{P}$, such that $\tilde{g}$ and $\tilde{P}$ verify the conditions (1.1) and (1.2). The structure $(\tilde{P}, \tilde{g})$ induces on the submanifold M a $(a, \epsilon)f$ Riemannian structure, which verifies the following properties:*

$$(2.3) \left\{ \begin{array}{l} (i) \quad P^2 X = \epsilon(X - \sum_{\alpha=1}^r u_{\alpha}(X) \xi_{\alpha}), \\ (ii) \quad u_{\alpha}(PX) = - \sum_{\beta=1}^r a_{\beta\alpha} u_{\beta}(X), \\ (iii) \quad a_{\alpha\beta} = \epsilon a_{\beta\alpha}, \\ (iv) \quad u_{\alpha}(\xi_{\beta}) = \delta_{\alpha\beta} - \epsilon \sum_{\gamma=1}^r a_{\alpha\gamma} a_{\gamma\beta}, \\ (v) \quad P\xi_{\alpha} = - \sum_{\beta=1}^r a_{\alpha\beta} \xi_{\beta}, \\ (vi) \quad u_{\alpha}(X) = g(X, \xi_{\alpha}), \\ (vii) \quad g(PX, Y) = \epsilon g(X, PY), \\ (viii) \quad g(PX, PY) = g(X, Y) - \\ \quad - \sum_{\alpha=1}^r u_{\alpha}(X) u_{\alpha}(Y) \end{array} \right.$$

for any $X, Y \in \chi(M)$ and $\alpha, \beta \in \{1, \dots, r\}$.

Remark 2.1. From (2.3)(i), (v), (vi) we obtain

$$(2.4) \quad \tilde{P}^3 X = \epsilon PX + \sum_{\alpha, \beta=1}^r a_{\alpha\beta} g(X, \xi_{\beta}) \xi_{\alpha}$$

for every $X \in \chi(M)$.

Let $\tilde{\nabla}$ and ∇ be the Levi-Civita connections defined on M and $\tilde{M}$ respectively, with respect to $\tilde{g}$ and g respectively.

Definition 2.2. If $(\tilde{M}, \tilde{g}, \tilde{P})$ is an almost product Riemannian manifold such that $\tilde{\nabla}\tilde{P} = 0$, then we say that $\tilde{M}$ is a locally product Riemannian manifold.

The Nijenhuis torsion tensor field of P has the form $N_P(X, Y) = [PX, PY] + P^2[X, Y] - P[PX, Y] - P[X, PY]$ for any $X, Y \in \chi(M)$. As in the case of an almost paracontact structure ([10]), one can defined the normal $(P, g, u_{\alpha}, \xi_{\alpha}, (a_{\alpha\beta})_r)$ structure on M .

Definition 2.3. If we have the equality

$$(2.5) \quad N_P(X, Y) - 2 \sum_{\alpha} du_{\alpha}(X, Y) \xi_{\alpha} = 0,$$

for any $X, Y \in \chi(M)$, then the $(a, 1)f$ induced structure on submanifold M in a Riemannian almost product manifold $(\tilde{M}, \tilde{g}, \tilde{P})$ is said to be normal.

Theorem 2.2. ([7]) Let M be an n -dimensional submanifold of codimension r in a locally product Riemannian manifold $(\tilde{M}, \tilde{g}, \tilde{P})$ and we suppose that the normal connection $\nabla^{\perp}$ vanishes identically. If the determinant of the matrix $(I_r - a^2)$ does not vanish in any point $x \in M$, then the $(a, 1)f$ induced structure on M is normal if and only if the $(1, 1)$ tensor field P commutes with the Weingarten operators A_{α} (for every $\alpha \in \{1, \dots, r\}$).

Remark 2.2. If the submanifold M in $\widetilde{M}$ is totally umbilical, then the commutativity between the $(1,1)$ tensor field P and the Weingarten operators A_α (for any $\alpha \in \{1, \dots, r\}$) has taken place. Therefore, if the determinant of the matrix $(I_r - a^2)$ does not vanish in any point $x \in M$ then the $(a, 1)f$ induced structure on a totally umbilical submanifold is normal.

3 Examples of $(a, 1)f$ Riemannian structures on spheres or product of spheres in Euclidean space

We suppose that the ambient space is E^{2p+q} ($p, q \in \mathbb{N}^*$) and for any point of E^{2p+q} we have its coordinates:

$$(x^1, \dots, x^p, y^1, \dots, y^p, z^1, \dots, z^q) := (x^i, y^i, z^j)$$

where $i \in \{1, \dots, p\}$ and $j \in \{1, \dots, q\}$. The tangent space $T_x(E^{2p+q})$ is isomorphic with E^{2p+q} .

Let $\tilde{P} : E^{2p+q} \rightarrow E^{2p+q}$ an almost product structure on E^{2p+q} such that

$$\begin{aligned} (3.1) \quad \tilde{P}(x^1, \dots, x^p, y^1, \dots, y^p, z^1, \dots, z^q) &= \\ &= (y^1, \dots, y^p, x^1, \dots, x^p, \varepsilon z^1, \dots, \varepsilon z^q) \end{aligned}$$

where $\varepsilon = \pm 1$. Thus, $(\tilde{P}, \langle \rangle)$ is an almost product Riemannian structure on E^{2p+q} .

Example 1. In this example, we construct an $(a, 1)f$ -structure on the sphere $S^{2p+q-1}(R) \hookrightarrow E^{2p+q}$. The equation of sphere $S^{2p+q-1}(R)$ is

$$(3.2) \quad \sum_{i=1}^p (x^i)^2 + \sum_{i=1}^p (y^i)^2 + \sum_{j=1}^q (z^j)^2 = R^2$$

where R is its radius and $(x^1, \dots, x^p, y^1, \dots, y^p, z^1, \dots, z^q)$ are the coordinates of any point of $S^{2p+q-1}(R)$. We use the following notations

$$(3.3) \quad \sum_{i=1}^p (x^i)^2 = r_1^2, \quad \sum_{i=1}^p (y^i)^2 = r_2^2, \quad \sum_{j=1}^q (z^j)^2 = r_3^2,$$

and $r_1^2 + r_2^2 = r^2$. Thus, we have $r^2 + r_3^2 = R^2$. We remark that

$$(3.4) \quad N_1 := \frac{1}{R}(x^i, y^i, z^j), \quad i \in 1, \dots, p, j \in \{1, \dots, q\}$$

is a unit normal vector field on sphere $S^{2p+q-1}(R)$ and

$$(3.5) \quad \tilde{P}N_1 = \frac{1}{R}(y^i, x^i, \varepsilon z^j)$$

For a tangent vector field X on $S^{2p+q-1}(R)$ we use the following notation

$$(3.6) \quad X = (X^1, \dots, X^p, Y^1, \dots, Y^p, Z^1, \dots, Z^q) := (X^i, Y^i, Z^j)$$

Hence we have

$$(3.7) \quad \sum_{i=1}^p x^i X^i + \sum_{i=1}^p y^i Y^i + \sum_{j=1}^q z^j Z^j = 0$$

If we decompose $\tilde{P}N_1$ in the tangential and normal components at sphere $S^{2p+q-1}(R)$, we obtain

$$(3.8) \quad \tilde{P}N_1 = \bar{\xi} + a_{11}N_1$$

where $a_{11} = \langle \tilde{P}N_1, N_1 \rangle$. If we use the following notation

$$(3.9) \quad \sigma = \sum_{i=1}^p x^i y^i$$

we obtain

$$(3.10) \quad a_{11} = \frac{2\sigma + \varepsilon r_3^2}{R^2}.$$

From (3.8) we obtain the tangential component of $\tilde{P}N_1$ at sphere $S^{2p+q-1}(R)$:

$$(3.11) \quad \bar{\xi} = \frac{1}{R}(y^i - \frac{2\sigma + \varepsilon r_3^2}{R^2}x^i, x^i - \frac{2\sigma + \varepsilon r_3^2}{R^2}y^i, \frac{\varepsilon r^2 - 2\sigma}{R^2}z^j)$$

for $i \in \{1, \dots, p\}$ and $j \in \{1, \dots, q\}$.

If we decompose $\tilde{P}X$ in the tangential and normal components at sphere S^{2p+q-1} (where X is a tangent vector field at $S^{2p+q-1}(R)$), we obtain

$$(3.12) \quad \tilde{P}X = \bar{P}X + \bar{u}(X)N_1$$

From $\bar{u}(X) = \langle X, \bar{\xi} \rangle$ and (3.7) we obtain

$$(3.13) \quad \bar{u}(X) = \frac{1}{R}[\sum_{i=1}^p (x^i Y^i + y^i X^i) + \varepsilon \sum_{j=1}^q z^j Z^j]$$

From $\bar{P}X = \tilde{P}X - \bar{u}(X)N_1$ we obtain

$$(3.14) \quad \bar{P}X = (Y^i - \frac{\bar{u}(X)}{R}x^i, X^i - \frac{\bar{u}(X)}{R}y^i, \varepsilon Z^j - \frac{\bar{u}(X)}{R}z^j)$$

where $X := (X^i, Y^i, Z^j)$ is a tangent vector at sphere in any point (x^i, y^i, z^j) and $\bar{u}(X)$ was defined in (3.13).

Therefore, from the relations (3.10), (3.11), (3.13), (3.14) we have a $(\bar{P}, \langle, \rangle, \bar{\xi}, \bar{u}, a_{11})$ induced structure by $\tilde{P}$ from E^{2p+q} on the sphere $S^{2p+q-1}(R)$ of codimension 1 in Euclidean space E^{2p+q} . We use the following notation $\bar{a} := a_{11}$. If $a_{11} \neq 1$ in every point of $S^{2p+q-1}(R)$, from the Theorem 2.2 we obtain that the structure $(\bar{P}, \langle, \rangle, \bar{\xi}, \bar{u}, \bar{a})$ is an $(\bar{a}, 1)f$ normal structure, because $S^{2p+q-1}(R)$ is a totally umbilical hypersurface in E^{2p+q} and from this we have that the tensor field P commutes with the Weingarten operator A .

Example 2. In this example, we construct an $(\hat{a}, 1)f$ -structure on the product of spheres $S^{2p-1}(r) \times S^{q-1}(r_3)$.

Let E^{2p+q} ($p, q \in \mathbb{N}^*$) be Euclidean space ($p, q \in \mathbb{N}^*$) endowed with an almost product Riemannian structure $\tilde{P}$ defined in (3.1). It is obvious that $E^{2p+q} = E^{2p} \times E^q$ and in each of spaces E^{2p} and E^q respectively, we can get a hypersphere $S^{2p-1}(r) = \{(x^1, \dots, x^p, y^1, \dots, y^p), \sum_{i=1}^p ((x^i)^2 + (y^i)^2) = r^2\}$ and $S^{q-1}(r_3) = \{(z^1, \dots, z^q), \sum_{j=1}^q (z^j)^2 = r_3^2\}$ respectively, where $r^2 + r_3^2 = R^2$. We construct (in the same manner like in [8] or [7]) the product manifold $S^{2p-1}(r) \times S^{q-1}(r_3)$. Any point of $S^{2p-1}(r) \times S^{q-1}(r_3)$ has the coordinates $(x^1, \dots, x^p, y^1, \dots, y^p, z^1, \dots, z^q) := (x^i, y^i, z^j)$ which verifies (3.3). Thus, $S^{2p-1}(r) \times S^{q-1}(r_3)$ is a submanifold of codimension 2 in E^{2p+q} . Furthermore, $S^{2p-1}(r) \times S^{q-1}(r_3)$ is a submanifold of codimension 1 in $S^{2p+q-1}(R)$. Therefore we have the successive imbedded

$$S^{2p-1}(r) \times S^{q-1}(r_3) \hookrightarrow S^{2p+q-1}(R) \hookrightarrow E^{2p+q}$$

The tangent space in a point (x^i, y^i, z^j) at the product of spheres $S^{2p-1}(r) \times S^{q-1}(r_3)$ is $T_{(x^1, \dots, x^p, y^1, \dots, y^p, \underbrace{0, \dots, 0}_q)} S^{2p-1}(r) \oplus T_{(\underbrace{0, \dots, 0}_{2p}, z^1, \dots, z^q)} S^{q-1}(r_3)$.

A vector $(X^1, \dots, X^p, Y^1, \dots, Y^p)$ from $T_{(x^1, \dots, x^p, y^1, \dots, y^p)} E^{2p}$ is tangent to $S^{2p-1}(r)$ if and only if

$$(3.15) \quad \sum_{i=1}^p x^i X^i + \sum_{i=1}^p y^i Y^i = 0$$

and it can be identified by $(X^1, \dots, X^p, Y^1, \dots, Y^p, \underbrace{0, \dots, 0}_q)$ from E^{2p+q} . A

vector $(Z^1, \dots, Z^q)$ from $T_{(z^1, \dots, z^q)} E^q$ is tangent to $S^{q-1}(r_3)$ if and only if

$$(3.16) \quad \sum_{j=1}^q z^j Z^j = 0$$

and it can be identified by $(\underbrace{0, \dots, 0}_{2p}, Z^1, \dots, Z^q)$ from E^{2p+q} . Consequently,

for any point $(x^i, y^i, z^j) \in S^{2p-1}(r) \times S^{q-1}(r_3)$ we have $(X^i, Y^i, Z^j) \in T_{(x^1, \dots, x^p, y^1, \dots, y^p, z^1, \dots, z^q)} (S^{2p-1}(r) \times S^{q-1}(r_3))$ if and only if the relations (3.15)

and (3.16) are satisfied. Furthermore, we remark that (X^i, Y^i, Z^j) is a tangent vector field at $S^{2p+q-1}(R)$ and from this it follows that

$$T_{(x^i, y^i, z^j)}(S^{2p-1}(r) \times S^{q-1}(r_3)) \subset T_{(x^i, y^i, z^j)}S^{2p+q-1}(R),$$

for any point $(x^i, y^i, z^j) \in S^{2p-1}(r) \times S^{q-1}(r_3)$.

The normal unit vector field N_1 at $S^{2p+q-1}(R)$ is also a normal vector field at $(S^{2p-1}(r) \times S^{q-1}(r_3))$, when it is considered in its points. We construct an unit vector field N_2 on S^{2p+q-1} , by the relation

$$(3.17) \quad N_2 = \frac{1}{R}(\frac{r_3}{r}x^i, \frac{r_3}{r}y^i, -\frac{r}{r_3}z^j)$$

We remark that N_2 is orthogonal on N_1 . From (3.15) and (3.16) we obtain that N_2 is orthogonal at $S^{2p-1}(r) \times S^{q-1}(r_3)$. Thus, (N_1, N_2) is an orthonormal basis in $T_{(x^i, y^i, z^j)}^\perp S^{2p-1}(r) \times S^{q-1}(r_3)$ in any point $(x^i, y^i, z^j) \in S^{2p-1}(r) \times S^{q-1}(r_3)$.

From the decomposed of $\tilde{P}N_k$ in tangential and normal components at $S^{2p-1}(r) \times S^{q-1}(r_3)$, we obtain

$$(3.18) \quad \tilde{P}N_\alpha = \hat{\xi}_\alpha + a_{\alpha 1}N_1 + a_{\alpha 2}N_2, \quad \alpha \in \{1, 2\}.$$

a_{11} was defined in (3.10). From (2.3)(iii) we have $a_{12} = a_{21}$ and from $a_{\alpha\beta} = \langle \tilde{P}N_\alpha, N_\beta \rangle$ ($\alpha, \beta \in \{1, 2\}$), we obtain

$$(3.19) \quad a_{12} = a_{21} = \frac{(2\sigma - \varepsilon r^2)r_3}{rR^2}$$

and

$$(3.20) \quad a_{22} = \frac{2\sigma r_3^2 + \varepsilon r^4}{r^2 R^2}$$

From (3.18), (3.19), (3.20) and (3.10) we obtain

$$(3.21) \quad \hat{\xi}_1 = \frac{1}{R}(y^i - \frac{2\sigma}{r^2}x^i, x^i - \frac{2\sigma}{r^2}y^i, 0).$$

and

$$(3.22) \quad \hat{\xi}_2 = \frac{r_3}{rR}(y^i - \frac{2\sigma}{r^2}x^i, x^i - \frac{2\sigma}{r^2}y^i, 0).$$

Therefore, the matrix $\hat{a} := (a_{\alpha\beta})_2$ is given by

$$(3.23) \quad \hat{a} := \begin{pmatrix} \frac{2\sigma + \varepsilon r_3^2}{R^2} & \frac{(2\sigma - \varepsilon r^2)r_3}{rR^2} \\ \frac{(2\sigma - \varepsilon r^2)r_3}{rR^2} & \frac{2\sigma r_3^2 + \varepsilon r^4}{r^2 R^2} \end{pmatrix}$$

From the decomposed of $\tilde{P}X$ in tangential and normal components at $S^{2p-1}(r) \times S^{q-1}(r_3)$ (where $X := (X^i, Y^i, Z^j)$ is a tangent vector field on $S^{2p-1}(r) \times S^{q-1}(r_3)$), we obtain

$$(3.24) \quad \tilde{P}X = \hat{P}X + \hat{u}_1(X)N_1 + \hat{u}_2(X)N_2.$$

We use the following notation

$$(3.25) \quad \tau := \sum_{i=1}^p (x^i Y^i + y^i X^i)$$

From $\hat{u}_\alpha(X) = \langle X, \hat{\xi}_\alpha \rangle$ (with $\alpha \in \{1, 2\}$) and (3.15) we obtain

$$(3.26) \quad \hat{u}_1(X) = \frac{\tau}{R}, \quad \hat{u}_2(X) = \frac{r_3 \tau}{rR}$$

for any tangent vector X on $S^{2p-1}(r) \times S^{q-1}(r_3)$ in a point $(x^i, y^i, z^i) \in S^{2p-1}(r) \times S^{q-1}(r_3)$. From (3.24) and (3.26), we obtain the tangent component $\hat{P}X$ of $\tilde{P}X$ on $S^{2p-1}(r) \times S^{q-1}(r_3)$, which is given by

$$(3.27) \quad \hat{P}X = (Y^i - \frac{\tau}{r^2} x^i, X^i - \frac{\tau}{r^2} y^i, \varepsilon Z^j)$$

Consequently, we obtain the $(\hat{P}, \langle, \rangle, \hat{\xi}_1, \hat{\xi}_2, \hat{u}_1, \hat{u}_2, \hat{a})$ induced structure on the product of spheres $S^{2p-1}(r) \times S^{q-1}(r_3)$ by the almost product Riemannian structure $(\hat{P}, \langle, \rangle)$ which is an $(\hat{a}, 1)f$ Riemannian structure induced on the submanifold $S^{2p-1}(r) \times S^{q-1}(r_3)$ of codimension 2 in Euclidean space E^{2p+q} .

Example 3. In this example, we construct an $(a, 1)f$ -structure on the product of spheres $S^{p-1}(r_1) \times S^{p-1}(r_2) \times S^{q-1}(r_3)$.

Let E^{2p+q} be the Euclidean space endowed with an almost product Riemannian structure $\tilde{P}$ which was defined in (3.1). We have $E^{2p+q} = E^p \times E^p \times E^q$ and in each of spaces E^p we can get a hypersphere

$$(3.28) \quad S^{p-1}(r_1) = \{(x^1, \dots, x^p), \sum_{i=1}^p (x^i)^2 = r_1^2\},$$

and

$$(3.29) \quad S^{p-1}(r_2) = \{(y^1, \dots, y^p), \sum_{i=1}^p (y^i)^2 = r_2^2\},$$

respectively, with the assumption that $r_1^2 + r_2^2 = r^2$.

We construct the product manifold $S^{p-1}(r_1) \times S^{p-1}(r_2) \times S^{q-1}(r_3)$.

Any point x of $S^{p-1}(r_1) \times S^{p-1}(r_2) \times S^{q-1}(r_3)$ has the coordinates

$$(x^1, \dots, x^p, y^1, \dots, y^p, z^1, \dots, z^q) := (x^i, y^i, z^j)$$

which verifies (3.3). Thus, $S^{p-1}(r_1) \times S^{p-1}(r_2) \times S^{q-1}(r_3)$ is a submanifold of codimension 3 in E^{2p+q} . Furthermore, $S^{p-1}(r_1) \times S^{p-1}(r_2) \times S^{q-1}(r_3)$ is a submanifold of codimension 1 in $S^{2p-1}(r) \times S^{q-1}(r_3)$ and a submanifold of codimension 2 in $S^{2p+q-1}(R)$. Therefore, we have the successive imbedded $S^{p-1}(r_1) \times S^{p-1}(r_2) \times S^{q-1}(r_3) \hookrightarrow S^{2p-1}(r) \times S^{q-1}(r_3) \hookrightarrow S^{2p+q-1}(R) \hookrightarrow E^{2p+q}$

The tangent space in a point $x := (x^i, y^i, z^j)$ at the product of spheres $S^{p-1}(r_1) \times S^{p-1}(r_2) \times S^{q-1}(r_3)$ is

$$T_{(x^i, 0, 0)} S^{p-1}(r_1) \oplus T_{(0, y^i, 0)} S^{p-1}(r_2) \oplus T_{(0, 0, z^j)} S^{q-1}(r_3)$$

where $(x^i, 0, 0)$ is a notation for $(x^1, \dots, x^p, \underbrace{0, \dots, 0}_{p+q})$, $(0, y^i, 0)$ is a notation for $(\underbrace{0, \dots, 0}_p, y^1, \dots, y^p, \underbrace{0, \dots, 0}_q)$ and $(0, 0, z^j)$ is a notation for $(\underbrace{0, \dots, 0}_{2p}, \underbrace{z^1, \dots, z^q}_{2p})$.

A vector $(X^1, \dots, X^p)$ from $T_{(x^1, \dots, x^p)} E^p$ is tangent to $S^{p-1}(r_1)$ if and only if

$$(3.30) \quad \sum_{i=1}^p x^i X^i = 0$$

and it can be identified by $(X^1, \dots, X^p, \underbrace{0, \dots, 0}_{p+q})$ from E^{2p+q} .

A vector $(Y^1, \dots, Y^p)$ from $T_{(y^1, \dots, y^p)} E^p$ is tangent to $S^{p-1}(r_2)$ if and only if

$$(3.31) \quad \sum_{i=1}^p y^i Y^i = 0$$

and it can be identified by $(\underbrace{0, \dots, 0}_p, Y^1, \dots, Y^p, \underbrace{0, \dots, 0}_q)$ from E^{2p+q} .

Consequently, for any point $(x^i, y^i, z^j) \in S^{p-1}(r_1) \times S^{p-1}(r_2) \times S^{q-1}(r_3)$ we have

$$(X^i, Y^i, Z^j) \in T_{(x^i, y^i, z^j)} (S^{p-1}(r_1) \times S^{p-1}(r_2) \times S^{q-1}(r_3))$$

if and only if the relations (3.16), (3.30) and (3.31) are satisfied. Furthermore, we remark that (X^i, Y^i, Z^j) is a tangent vector field at the product of spheres $S^{2p-1}(r) \times S^{q-1}(r_3)$ and at $S^{2p+q-1}(R)$, respectively. Thus, it follows that

$$T_{(x^i, y^i, z^j)} (S^{p-1}(r_1) \times S^{p-1}(r_2) \times S^{q-1}(r_3)) \subset T_{(x^i, y^i, z^j)} (S^{2p-1}(r) \times S^{q-1}(r_3)),$$

for any point $(x^i, y^i, z^j) \in S^{p-1}(r_1) \times S^{p-1}(r_2) \times S^{q-1}(r_3)$.

The normal unit vector fields N_1 and N_2 at $(S^{2p-1}(r) \times S^{q-1}(r_3))$ are also normal vector fields at $(S^{p-1}(r_1) \times (S^{p-1}(r_2) \times S^{q-1}(r_3)))$, when they are considered in its points. We construct a unit vector field N_3 by the relation

$$(3.32) \quad N_3 = \frac{1}{r} \left(\frac{r_2}{r_1} x^i, -\frac{r_1}{r_2} y^i, 0 \right)$$

We remark that N_3 is orthogonal on N_1 and N_2 , respectively. Furthermore, from (3.16), (3.30) and (3.31) we obtain that N_3 is orthogonal at $S^{p-1}(r_1) \times S^{p-1}(r_1) \times S^{q-1}(r_3)$. Thus, (N_1, N_2, N_3) is an orthonormal basis in $T_{(x^i, y^i, z^j)}^\perp S^{p-1}(r_1) \times S^{p-1}(r_2) \times S^{q-1}(r_3)$ in any point $(x^i, y^i, z^j) \in S^{p-1}(r_1) \times S^{p-1}(r_2) \times S^{q-1}(r_3)$.

From the decomposed of $\tilde{P}N_\alpha$ (with $\alpha \in \{1, 2, 3\}$) in the tangential and normal components at $S^{p-1}(r_1) \times S^{p-1}(r_2) \times S^{q-1}(r_3)$, we obtain

$$(3.33) \quad \tilde{P}N_\alpha = \xi_\alpha + a_{\alpha 1}N_1 + a_{\alpha 2}N_2 + a_{\alpha 3}N_3,$$

for every $\alpha \in \{1, 2, 3\}$. From (3.10) we have a_{11} , from (3.19) we have $a_{12} = a_{21}$ and from (3.20) we have a_{22} . Furthermore, from (2.3)(iii) we have that $a_{13} = a_{31}$ and $a_{23} = a_{32}$.

It is obvious that

$$(3.34) \quad a_{\alpha\beta} = \langle \tilde{P}N_\alpha, N_\beta \rangle, \quad (\alpha, \beta \in \{1, 2, 3\}).$$

From this, we have

$$(3.35) \quad a_{13} = a_{31} = \frac{r_2^2 - r_1^2}{r_1 r_2 r R} \sigma$$

$$(3.36) \quad a_{23} = a_{32} = \frac{(r_2^2 - r_1^2)r_3}{r_1 r_2 r^2 R} \sigma$$

and

$$(3.37) \quad a_{33} = -\frac{2\sigma}{r^2}$$

From (3.33) we obtain

$$(3.38) \quad \xi_1 = \frac{1}{R} \left(y^i - \frac{\sigma}{r_1^2} x^i, x^i - \frac{\sigma}{r_2^2} y^i, 0 \right).$$

$$(3.39) \quad \xi_2 = \frac{r_3}{rR} \left(y^i - \frac{\sigma}{r_1^2} x^i, x^i - \frac{\sigma}{r_2^2} y^i, 0 \right).$$

and

$$(3.40) \quad \xi_3 = \frac{1}{r} \left(\frac{\sigma}{r_1 r_2} x^i - \frac{r_1}{r_2} y^i, \frac{r_2}{r_1} x^i - \frac{\sigma}{r_1 r_2} y^i, 0 \right).$$

Therefore, the matrix $a := (a_{\alpha\beta})_3$ is given by

$$(3.41) \quad a = \begin{pmatrix} \frac{2\sigma + \varepsilon r_3^2}{R^2} & \frac{(2\sigma - \varepsilon r^2)r_3}{rR^2} & \frac{r_2^2 - r_1^2}{r_1 r_2 r R} \sigma \\ \frac{(2\sigma - \varepsilon r^2)r_3}{rR^2} & \frac{2\sigma r_3^2 + \varepsilon r^4}{r^2 R^2} & \frac{(r_2^2 - r_1^2)r_3}{r_1 r_2 r^2 R} \sigma \\ \frac{r_2^2 - r_1^2}{r_1 r_2 r R} \sigma & \frac{(r_2^2 - r_1^2)r_3}{r_1 r_2 r^2 R} \sigma & -\frac{2\sigma}{r^2} \end{pmatrix}$$

From the decomposed of $\tilde{P}X$ in tangential and normal components at the product of spheres $S^{p-1}(r_1) \times S^{p-1}(r_2) \times S^{q-1}(r_3)$ (where $X := (X^i, Y^i, Z^j)$ is a tangent vector field on $S^{p-1}(r_1) \times S^{p-1}(r_2) \times S^{q-1}(r_3)$), we obtain

$$(3.42) \quad \tilde{P}X = PX + u_1(X)N_1 + u_2(X)N_2 + u_3(X)N_3.$$

From (2.3)(vi) we have $u_\alpha(X) = \langle X, \xi_\alpha \rangle$ (with $\alpha \in \{1, 2, 3\}$) and using (3.37) and (3.38) we obtain

$$(3.43) \quad u_1(X) = \frac{\tau}{R},$$

$$(3.44) \quad u_2(X) = \frac{r_3\tau}{rR}$$

and respectively

$$(3.45) \quad u_3(X) = \frac{1}{r} \left[\frac{r_2}{r_1} \sum_{i=1}^p x^i Y^i - \frac{r_1}{r_2} \sum_{i=1}^p y^i X^i \right].$$

for any tangent vector $X := (X^i, Y^i, Z^j)$ in any point (x^i, y^i, z^j) on the product of spheres $S^{p-1}(r_1) \times S^{p-1}(r_2) \times S^{q-1}(r_3)$, with the notation (3.25) for τ . From (3.42), (3.43), (3.44) and (3.45), we obtain the tangent component PX of $\tilde{P}X$ in any point (x^i, y^i, z^j) on $S^{p-1}(r_1) \times S^{p-1}(r_2) \times S^{q-1}(r_3)$, which has the form

$$(3.46) \quad PX = (Y^i - \frac{\sum_{k=1}^p x^k Y^k}{r_1^2} x^i, X^i - \frac{\sum_{k=1}^p y^k X^k}{r_2^2} y^i, \varepsilon Z^j)$$

Consequently, we obtain the induced structure $(P, \langle, \rangle, \xi_1, \xi_2, \xi_3, u_1, u_2, u_3, (a_{\alpha\beta})_3)$ on $S^{p-1}(r_1) \times S^{p-1}(r_2) \times S^{q-1}(r_3)$ by the almost product Riemannian structure $(\tilde{P}, \langle, \rangle)$, which is an $(a, 1)f$ Riemannian structure induced on the product of spheres $S^{p-1}(r_1) \times S^{p-1}(r_2) \times S^{q-1}(r_3)$ in E^{2p+q} .

Remark 3.1. *If we consider the almost product structure on E^{2p+q} given by $\tilde{P} : E^{2p+q} \rightarrow E^{2p+q}$,*

$$\begin{aligned} \tilde{P}(x^1, \dots, x^p, y^1, \dots, y^p, z^1, \dots, z^q) = \\ = (y^1, \dots, y^p, x^1, \dots, x^p, \varepsilon_1 z^1, \dots, \varepsilon_q z^q) \end{aligned}$$

where $\varepsilon_j = \pm 1$ (for every $j \in \{1, \dots, q\}$), we obtain a generalization of $(a, 1)f$ induced structures defined in examples from above.

References

- [1] T. Adati, *Submanifolds of an almost product Riemannian manifold*, *Kodai Math. J.*, 4-2 (1981), 327-343.
- [2] D.E.Blair, G.D. Ludden, K. Yano, *Induced structures on submanifolds*, *Kodai Math. Sem. Rep.* 22 (1970), 188-198.
- [3] A. Bucki, A. Miernowski, *Invariant hypersurfaces of an lmost r-paracontact manifold*, *Demonstr. Math.*, 19(1), (1986), 113-121.
- [4] B.Y. Chen, *Geometry of Submanifolds*, Marcel Dekker Inc, New York (1973).
- [5] C.E. Hreţcanu, *On submanifolds in Riemannian almost product manifolds*, *MASEE International Congress on Mathematics MICOM 2006*, Cyprus
- [6] C.E. Hreţcanu, *A structure of paracontact type on submanifolds in almost product manifolds*, 5 th *Symposium on Mathematics Applied in Biology and Biophysics*, 2006, U.A.S.V.M Iasi, Romania.
- [7] C.E. Hreţcanu, *Induced structures on submanifolds in almost product Riemannian manifolds*, *arXiv:math.DG/0608533 v1 22 Aug 2006*.
- [8] S. Ianuş, *Differential geometry with applications to general relativity*, Editura Academiei RSR, Bucureşti, 1983.
- [9] K. Matsumoto, *On Submanifolds of Locally Product Riemannian Manifolds*, *TRU Mathematics* 18-2, 1982, 145-157.
- [10] I. Mihai, R. Roşca, L. Verstraelen, *Some aspects of the differential geometry of vector fields*, *Katholieke Universiteit Leuven*, PADGE, vol 2, 1996.
- [11] T. Miyazawa, *Hypersurfaces immersed in an almost product Riemannian manifold*, *Tensor N. S.*, 33-1 (1979), 114-116.
- [12] J. Nikic, *Conditions for invariant submanifold of a manifold with the (ϕ, ξ, η, G) -structure*, *Kragujevac J. Math.* 25 (2003), pag 147-154.
- [13] G. Pitiş, *On some submanifolds of a locally product manifold*, *Kodai Math.J.* 9 (1986), 327333
- [14] I. Sato, *On a structure similar to the almost contact structure I;II*, *Tensor N.S.* 30 (1976), 219-224; 31 (1977), 199-205.
- [15] X. Senlin and N. Yilong, *Submanifolds of Product Riemannian Manifold*, *Acta Mathematica Scientia* 2000, 20(B) 213-218.

- [16] K. Yano, M. Kon, *Structures on Manifolds*, World Scientific, Singapore, *Series in pure mathematics - Volume 3*, 1984.
- [17] K. Yano, M. Okumura, *On normal (f, g, u, v, λ) -structures on submanifolds of codimension 2 in an even-dimensional Euclidean space*, *Kodai Math. Sem. Rep* 23 (1971), 172-197.
- [18] K. Yano, M. Okumura, *On (f, g, u, v, λ) -structures*, *Kodai Math. Sem. Rep.* 22 (1970), 401-423.

"Ștefan cel Mare" University, Suceava, Romania
 E-mail address: cristinah@usv.ro